

GenCore version 5.1.4_p5_4578
Copyright (c) 1993 - 2003 CompuGen Ltd.

OM protein - protein search, using sw model

Run on: March 13, 2003, 08:31:55 ; Search time 13 Seconds
(without alignments)
79.762 Million cell updates/sec

Title: US-09-913-524-9

Perfect score: 143

Sequence: 1 PWSPALRLQRPPEPSAHAFCHR 25

Scoring table:

BLOSUM62
Gapop 10.0 , Gapext 0.5

Searched: 112892 seqs, 41476328 residues

Total number of hits satisfying chosen parameters: 112892

Minimum DB seq length: 0

Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%

Maximum Match 100%

Listing first 45 summaries

Database : SwissProt_40:*

Pred. No. is the number of results predicted by chance to have a
score greater than or equal to the score of the result being printed,
and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Score	Query Match %	Length	DB ID	Description
1	143	100.0	366	1 IHA_RAT	P17490 rattus norv
2	140	97.9	366	1 IHA_MOUSE	O04957 mus musculu
3	131	91.6	265	1 IHA_SHEEP	P38440 ovis aries
4	131	91.6	360	1 IHA_BOVIN	P07994 bos taurus
5	131	91.6	367	1 IHA_HORSE	P55101 equus cabal
6	128	89.5	366	1 IHA_HUMAN	P05111 homo sapien
7	127	88.8	364	1 IHA_PIG	P04087 sus scrofa
8	120	83.9	361	1 IHA_TRITU	O77555 trichosurus
9	93	65.0	329	1 IHA_CHICK	P43031 gallus gall
10	54	37.8	2504	1 FAS_HUMAN	P49327 homo sapien
11	52.5	36.7	368	1 GP62_HUMAN	Q9b217 homo sapien
12	51	35.7	451	1 TBG1_HUMAN	P23258 homo sapien
13	51	35.7	451	1 TBG1_MOUSE	Q92310 mus musculu
14	51	35.7	451	1 TBG2_HUMAN	Q9a133 homo sapien
15	51	35.7	451	1 TBG2_MOUSE	Q9bck3 mus musculu
16	51	35.7	451	1 TBG_XENIA	P23330 xenopus lae
17	50	35.0	228	1 Z113_MOUSE	Q9et26 mus musculu
18	49	34.3	224	1 TIM4_MOUSE	Q9jhb3 mus musculu
19	49	34.3	228	1 Z113_HUMAN	O9y508 homo sapien
20	49	34.3	352	1 TPO_CANFA	P42705 canis famil
21	48.5	33.9	646	1 PIXB_HUMAN	Q14155 homo sapien
22	48.5	33.9	646	1 PIXB_RAT	O55043 rattus norv
23	48	33.6	192	1 NIF2_FRAAL	P46040 frankia aln
24	47.5	33.2	886	1 SM6B_MOUSE	O54951 mus musculu
25	47.5	33.2	887	1 SM6B_RAT	O70141 rattus norv
26	47	32.9	528	1 DRTS_DAUCA	P45350 daucus caro
27	47	32.9	622	1 BIN4_MOUSE	Q9z0h1 mus musculu
28	47	32.9	1122	1 HDAS_HUMAN	Q9uq16 homo sapien
29	47	32.9	1428	1 ATRN_MOUSE	O9wu60 mus musculu
30	47	32.9	1429	1 ATRN_HUMAN	O75882 homo sapien
31	47	32.9	1493	1 M3K1_MOUSE	P53349 mus musculu
32	46.5	32.5	391	1 NIFP_PSEST	O52521 pseudomonas
33	46.5	32.5	428	1 SYH_CHLMU	Q9pfj9 chlamydia m

RESULT 1

ID	IHA_RAT	STANDARD	PRT	366 AA
AC	P17490			
DT	01-AUG-1990 (Rel. 15, Created)			
DT	01-AUG-1990 (Rel. 15, Last sequence update)			
DT	15-JUN-2002 (Rel. 41, Last annotation update)			
DE	Inhibin alpha chain precursor.			
GN	INHA.			
OS	Rattus norvegicus (Rat).			
OC	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;			
OC	Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Rattus.			
OX	NCBI_TaxID=10116;			
RN	[1]			
RP	SEQUENCE FROM N.A.			
RC	TISSUE=Liver;			
RX	MEDLINE=90190649; PubMed=2628729;			
RA	Peng Z.-M., Li Y.-P., Chen C.-L.C.;			
RT	"Analysis of the 5'-flanking regions of rat inhibin alpha- and			
RT	beta-B-subunit genes suggests two different regulatory mechanisms.";			
RL	Mol. Endocrinol. 3:1914-1925(1989).			
RN	[2]			
RP	SEQUENCE FROM N.A.			
RX	MEDLINE=91042598; PubMed=3153478;			
RA	Woodruff T.K., Meunier H., Jones P.B.C., Hsueh A.J.W., Mayo K.E.;			
RT	"Rat inhibin: molecular cloning of alpha- and beta-subunit			
RL	complementary deoxyribonucleic acids and expression in the ovary.";			
RL	Mol. Endocrinol. 1:561-568(1987).			
RN	[3]			
RP	SEQUENCE FROM N.A.			
RX	MEDLINE=90331931; PubMed 2484214;			
RA	Esch F.S., Shimada S., Cocksey K., Mercado M., Mason A.J.,			
RA	Ying S.Y., Ueno N., Ling N.;			
RT	"Complementary deoxyribonucleic acid (cDNA) cloning and DNA sequence			
RL	analysis of rat ovarian inhibins.";			
RL	Mol. Endocrinol. 1:388-396(1987).			
RN	[4]			
RP	TISSUE SPECIFICITY.			
RX	MEDLINE=88124801; PubMed=2829170;			
RA	Meunier H., Rivier C., Evans R.M., Vale W.;			
RT	"Gonadal and extragonadal expression of inhibin alpha, beta A, and			
RT	beta B subunits in various tissues predicts diverse functions.";			
CC	Proc. Natl. Acad. Sci. U.S.A. 85:247-251(1988).			
CC	-!- FUNCTION: INHIBINS AND ACTIVINS INHIBIT AND ACTIVATE,			
CC	RESPECTIVELY, THE SECRETION OF FOLLITROPIN BY THE PITUITARY GLAND.			
CC	INHIBINS/ACTIVINS ARE INVOLVED IN REGULATING A NUMBER OF DIVERSE			
CC	FUNCTIONS SUCH AS HYPOTHALAMIC AND PITUITARY HORMONE SECRETION,			
CC	GONADAL HORMONE SECRETION, GERM CELL DEVELOPMENT AND MATURATION,			
CC	ERYTHROID DIFFERENTIATION, INSULIN SECRETION, NERVE CELL SURVIVAL,			
CC	EMBRYONIC AXIAL DEVELOPMENT OR BONE GROWTH, DEPENDING ON THEIR			
CC	SUBUNIT COMPOSITION. INHIBINS APPEAR TO OPPOSE THE FUNCTIONS OF			
CC	ACTIVINS.			
CC	-!- SUBUNIT: DIMERIC, LINKED BY ONE OR MORE DISULFIDE BONDS.			
CC	INHIBIN A IS A DIMER OF ALPHA AND BETA-A.			
CC	INHIBIN B IS A DIMER OF ALPHA AND BETA-B.			
CC	-!- TISSUE SPECIFICITY: MAINLY EXPRESSED IN OVARY AND TESTIS. ALPHA-			

P13100 pneumocysti
O15213 homo sapien
P23799 trypanosoma
P10358 turnip yell
P20128 turnip yell
P28477 turnip yell
Q9es28 mus musculu
Q9h3t3 homo sapien
P19711 bovine vira
Q61337 mus musculu
O35147 rattus norv
P45351 cryptococcu

[3] SEQUENCE OF 49-366 FROM N.A.
RP STRAIN-Swiss Webster;
RC MEDLINE-91071531; PubMed=2253839;
RX Tone S., Katoh Y., Fujimoto H., Togashi S., Yanazawa M., Kato Y.,
RA Higashinakagawa T.,
RT "Expression of inhibin alpha-subunit gene during mouse
RT gametogenesis.";
RL Differentiation 44:62-68(1990).
RN [4]
RP FUNCTION.
RX MEDLINE-93078845; PubMed=1448148;
RA Matzuk M.M., Finegold M.J., Su J.G.W., Hsueh A.J.W., Bradley A.;
RT "Alpha-inhibin is a tumour-suppressor gene with gonadal specificity in
RT mice.";
RL Nature 360:313-319(1992).
RN [5]
RP FUNCTION.
RX MEDLINE-94377444; PubMed=8090730;
RA Matzuk M.M., Finegold M.J., Mather J.P., Krummen L., Lu H.,
RA Bradley A.;
RT "Development of cancer cachexia-like syndrome and adrenal tumors in
RT inhibin-deficient mice.";
RL Proc. Natl. Acad. Sci. U.S.A. 91:8817-8821(1994).
RN [6]
RP FUNCTION, AND REVIEW.
RX MEDLINE-96321780; PubMed=8701077;
RA Matzuk M.M., Kumar T.R., Shou W., Coerver K.A., Lau A.L.,
RA Behringer R.R., Finegold M.J.;
RT "Transgenic models to study the roles of inhibins and activins in
RT reproduction, oncogenesis, and development.";
RL Recent Prog. Horm. Res. 51:123-154(1996).
RN -!- FUNCTION: INHIBINS AND ACTIVINS INHIBIT AND ACTIVATE,
CC RESPECTIVELY, THE SECRETION OF FOLLITROPIN BY THE PITUITARY GLAND.
CC INHIBINS/ACTIVINS ARE INVOLVED IN REGULATING A NUMBER OF DIVERSE
CC FUNCTIONS SUCH AS HYPOTHALAMIC AND PITUITARY HORMONE SECRETION,
CC GONADAL HORMONE SECRETION, GERM CELL DEVELOPMENT AND MATURATION,
CC ERYTHROID DIFFERENTIATION, INSULIN SECRETION, NERVE CELL SURVIVAL,
CC EMBRYONIC AXIAL DEVELOPMENT OR BONE GROWTH, DEPENDING ON THEIR
CC SUBUNIT COMPOSITION. INHIBINS APPEAR TO OPPOSE THE FUNCTIONS OF
CC ACTIVINS. INHIBIN DEFICIENT MICE ARE VIABLE BUT ARE ACUTELY
CC SENSITIVE TO DEVELOPMENT OF GONADAL SEX-CORD STROMAL TUMORS.
RN -!- SUBUNIT: DIMERIC, LINKED BY ONE OR MORE DISULFIDE BONDS.
CC INHIBIN A IS A DIMER OF ALPHA AND BETA-A.
CC INHIBIN B IS A DIMER OF ALPHA AND BETA-B.
RN -!- SIMILARITY: BELONGS TO THE TGF-BETA FAMILY.
CC -----
CC This SWISS-PROT entry is copyright. It is produced through a collaboration
CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC the European Bioinformatics Institute. There are no restrictions on its
CC use by non-profit institutions as long as its content is in no way
CC modified and this statement is not removed. Usage by and for commercial
CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
CC or send an email to license@isb-sib.ch).
CC -----
CC EMBL; X69618; CAA49324.1; -
DR EMBL; M95525; AAA39314.1; -
DR EMBL; M95526; AAA39314.1; JOINED.
DR EMBL; X55957; CAA39424.1; -
DR PIR; S31439; S31439.
DR PIR; JC1106; JC1106.
DR HSSP; P18075; IBMP.
DR MGD; MGI:96569; Inha.
DR InterPro; IPR002405; Inhibin_alpha.
DR InterPro; IPR001839; TGFb.
DR Pfam; PF00019; TGF-beta; 1.
DR PRINTS; PR00669; INHIBINA.
DR ProDom; PD000357; TGFb; 1.
DR SMART; SW00204; TGFb; 1.
DR PROSITE; PS00250; TGF_BETA_1; 1.
KW Growth factor; Hormone; Glycoprotein; Signal.
FT SIGNAL 1 20
FT PROPEL 21 233
FT POTENTIAL.

```

FT CHAIN 234 366 INHIBIN ALPHA CHAIN.
FT DISULFID 263 328 BY SIMILARITY.
FT DISULFID 292 363 BY SIMILARITY.
FT DISULFID 296 365 BY SIMILARITY.
FT DISULFID 327 327 INTERCHAIN (BY SIMILARITY).
FT CARBOHYD 147 147 N-LINKED (GLCNAC. . .) (POTENTIAL).
FT CARBOHYD 269 269 N-LINKED (GLCNAC. . .) (POTENTIAL).
FT CONFLICT 165 165 A -> R (IN REF. 2).
FT CONFLICT 171 171 V -> G (IN REF. 2 AND 3).
FT CONFLICT 336 336 T -> R (IN REF. 2 AND 3).
SQ SEQUENCE 366 AA; 39536 MW; 8F3851B722FE0011 CRC64;

Query Match 97.9%; Score 140; DB 1; Length 366;
Best Local Similarity 96.0%; Pred. No. 2.5e-12;
Matches 24; Conservative 1; Mismatches 0; Indels 0; Gaps 0;

QY 1 PWSPAALRLQRPPEPSAHAFCHR 25
Db 241 PWSPAALRLQRPPEPSAHAFCHR 265
|||||
|||||

RESULT 3
IHA_SHEEP
ID IHA_SHEEP STANDARD; PRT; 265 AA.
AC P38440;
DT 01-OCT-1994 (Rel. 30, Created)
DT 01-OCT-1994 (Rel. 30, Last sequence update)
DT 16-OCT-2001 (Rel. 40, Last annotation update)
DE Inhibin alpha chain (fragment).
GN INHA.
OS Ovis aries (Sheep).
OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovidae;
OC Bovidae; Caprinae; Ovis.
OX NCBI_TaxID=9940;
RN 1;
RP SEQUENCE FROM N.A.
RC Tissue-Ovary;
RA Sadanandan S.L., Jeyaseelan K.;
RL Submitted (FEB-1994) to the EMBL/GenBank/DBJ databases.
-!- FUNCTION: INHIBINS AND ACTIVINS INHIBIT AND ACTIVATE,
RESPECTIVELY. THE SECRETION OF FOLLITROPIN BY THE PITUITARY GLAND.
INHIBIN A IS A DIMER OF ALPHA AND BETA-A.
INHIBIN B IS A DIMER OF ALPHA AND BETA-B.
-!- SIMILARITY: BELONGS TO THE TGF-BETA FAMILY.
-----
CC This SWISS-PROT entry is copyright. It is produced through a collaboration
CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC the European Bioinformatics Institute. There are no restrictions on its
CC use by non-profit institutions as long as its content is in no way
CC modified and this statement is not removed. Usage by and for commercial
CC entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC or send an email to license@isb-sib.ch).
-----
CC EMBL; I28815; AAA31553.1; -
CC InterPro; IPR001839; TGFb.
CC Pfam; PF00019; TGF-beta; 1.
CC ProDom; PD000357; TGFb; 1.
CC SMART; SM00204; TGFb; 1.
CC PROSITE; PS00250; TGF_BETA_1; 1.
KW Growth factor; Hormone; Glycoprotein.
FT NON_TER 1
FT DISULFID 161 227 BY SIMILARITY.
FT DISULFID 190 262 BY SIMILARITY.
FT DISULFID 194 264 BY SIMILARITY.

```

```

FT DISULFID 226 226 INTERCHAIN (BY SIMILARITY).
FT CARBOHYD 45 45 N-LINKED (GLCNAC. . .) (POTENTIAL).
FT CARBOHYD 167 167 N-LINKED (GLCNAC. . .) (POTENTIAL).
SQ SEQUENCE 265 AA; 28754 MW; D880E9AB156B656 CRC64;

Query Match 91.6%; Score 131; DB 1; Length 265;
Best Local Similarity 92.0%; Pred. No. 3.2e-11;
Matches 23; Conservative 1; Mismatches 1; Indels 0; Gaps 0;

QY 1 PWSPAALRLQRPPEPSAHAFCHR 25
Db 139 PWSPAALRLQRPPEPSAHAFCHR 163
|||||
|||||

RESULT 4
IHA_BOVIN
ID IHA_BOVIN STANDARD; PRT; 360 AA.
AC P07994;
DT 01-AUG-1988 (Rel. 08, Created)
DT 01-AUG-1988 (Rel. 08, Last sequence update)
DT 16-OCT-2001 (Rel. 40, Last annotation update)
DE Inhibin alpha chain precursor.
GN INHA.
OS Bos taurus (Bovine).
OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovidae;
OC Bovidae; Bovinae; Bos.
OX NCBI_TaxID=9913;
RN 1;
RP SEQUENCE FROM N.A.
RC Tissue-Follicular fluid;
RX MEDLINE=86205842; PubMed=3458167;
RA Forage R.G., Ring J.M., Brown K.W., McInerney B.V., Cobon G.S.,
RA Gresson R.P., Robertson D.M., Morgan F.J., Hearn M.T.W., Findlay J.K.,
RA Wettienhall R.E.H., Burger H.G., de Kretser D.M.;
RT "Cloning and sequence analysis of cDNA species coding for the two
RT subunits of inhibin from bovine follicular fluid.";
RL Proc. Natl. Acad. Sci. U.S.A. 83:3091-3095(1986).
RN 12;
RP SEQUENCE OF 1-87 FROM N.A.
RC Tissue-Liver;
RX MEDLINE=95112839; PubMed=7813485;
RA Thompson D.A., Grouin C.N., Martin F.;
RT "Genomic cloning and sequence analyses of the bovine alpha-, beta A-
RT and beta B-inhibin/activin genes. Identification of transcription
RT factor AP-2-binding sites in the 5'-flanking regions by DNase I
RT footprinting.";
RL Eur. J. Biochem. 226:751-764(1994).
-!- FUNCTION: INHIBINS AND ACTIVINS INHIBIT AND ACTIVATE,
RESPECTIVELY. THE SECRETION OF FOLLITROPIN BY THE PITUITARY GLAND.
INHIBINS/ACTIVINS ARE INVOLVED IN REGULATING A NUMBER OF DIVERSE
CC FUNCTIONS SUCH AS HYPOPHALAMIC AND PITUITARY HORMONE SECRETION,
CC GONADAL HORMONE SECRETION, GERM CELL DEVELOPMENT AND MATURATION,
CC ERYTHROID DIFFERENTIATION, INSULIN SECRETION, NERVE CELL SURVIVAL,
CC EMBRYONIC AXIAL DEVELOPMENT OR BONE GROWTH, DEPENDING ON THEIR
CC SUBUNIT COMPOSITION. INHIBINS APPEAR TO OPPOSE THE FUNCTIONS OF
CC ACTIVINS.
-!- SUBUNIT: DIMERIC, LINKED BY ONE OR MORE DISULFIDE BONDS.
INHIBIN A IS A DIMER OF ALPHA AND BETA-A.
INHIBIN B IS A DIMER OF ALPHA AND BETA-B.
-!- SIMILARITY: BELONGS TO THE TGF-BETA FAMILY.
-----
CC This SWISS-PROT entry is copyright. It is produced through a collaboration
CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC the European Bioinformatics Institute. There are no restrictions on its
CC use by non-profit institutions as long as its content is in no way
CC modified and this statement is not removed. Usage by and for commercial
CC entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC or send an email to license@isb-sib.ch).
-----
CC EMBL; M13273; AAA97414.1; -
CC EMBL; A14416; CAA01156.1; -
CC EMBL; U16237; AAB60262.1; -

```

```

DR PR: A25732; A25732.
DR InterPro: IPR002405; Inhibin_alpha.
DR InterPro: IPR001839; TGFb.
DR Pfam: PF00019; TGF-beta; 1.
DR PRINTS: PR00669; INHIBIN.
DR PRODOM: PD000357; TGFb; 1.
DR SMART: SM00204; TGFb; 1.
DR PROSITE: PS00250; TGF_BETA_1; 1.
KW Growth factor; Hormone; Glycoprotein; Signal.
FT SIGNAL 1 17 POTENTIAL.
FT PROPEP 18 226
FT CHAIN 227 360 INHIBIN ALPHA CHAIN.
FT DISULFID 236 322 BY SIMILARITY.
FT DISULFID 285 357 BY SIMILARITY.
FT DISULFID 289 359 BY SIMILARITY.
FT DISULFID 321 321 INTERCHAIN (BY SIMILARITY).
FT CARBOHYD 140 140 N-LINKED (GLCNAC. . .) (POTENTIAL).
FT CARBOHYD 262 262 N-LINKED (GLCNAC. . .) (POTENTIAL).
SQ SEQUENCE 360 AA; 38409 MW; FBF385DDIEFE46 CRC64;

Query Match 91.68; Score 131; DB 1; Length 360;
Best Local Similarity 92.08; Pred. No. 4.4e-11;
Matches 23; Conservative 1; Mismatches 1; Indels 0; Gaps 0;

QY 1 PWSPAALRLIQRPPEPSAHAFCHR 25
| | | | | | | | | | | | | | | | | | | |
DB 234 PWSPAALRLIQRPPEPSAHADCHR 258

RESULT 5
IHA_HORSE
ID IHA_HORSE STANDARD; PRI; 367 AA.
AC P55101; Q28470;
DT 01-OCT-1996 (Rel. 34, Created)
DT 16-OCT-2001 (Rel. 40, Last sequence update)
DE Inhibin alpha chain precursor.
GN INHA.
OS Equus caballus (Horse).
OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC Mammalia; Theriata; Perissodactyla; Equidae; Equus.
OX NCBI_TaxId:9796;
RN [1]
RP SEQUENCE FROM N.A.
RC TISSUE-Ovary.
RX MEDLINE:96155857; PubMed:8593300;
RA Yamaguchi K., Yoshida S., Hasegawa T., Ikeda A., Chang K.T.,
RA Matsuyama S., Nishihara M., Miyazawa K., Takahashi M.;
RT "Molecular cloning of DNA for inhibin alpha-subunit from equine
RT ovary.";
RL J. Vet. Med. Sci. 57:905-909(1995).
RN [2]
RP SEQUENCE OF 98-367 FROM N.A.
RC TISSUE-Testis;
RA Adams M.H., Baker C.B., McDowell K.J.;
RT "Molecular cloning and sequencing of equine inhibin alpha cDNA.";
RL Anim. Biotechnol. 7:1-9(1996).
CC -!- FUNCTION: INHIBINS AND ACTIVINS INHIBIT AND ACTIVATE,
CC RESPECTIVELY, THE SECRETION OF FOLLITROPIN BY THE PITUITARY GLAND.
CC INHIBINS/ACTIVINS ARE INVOLVED IN REGULATING A NUMBER OF DIVERSE
CC FUNCTIONS SUCH AS HYPOTHALAMIC AND PITUITARY HORMONE SECRETION,
CC GONADAL HORMONE SECRETION, GERM CELL DEVELOPMENT AND MATURATION,
CC ERYTHROID DIFFERENTIATION, INSULIN SECRETION, NERVE CELL SURVIVAL,
CC EMBRYONIC AXIAL DEVELOPMENT OR BONE GROWTH, DEPENDING ON THEIR
CC SUBUNIT COMPOSITION. INHIBINS APPEAR TO OPPOSE THE FUNCTIONS OF
CC ACTIVINS.
CC -!- SUBUNIT: DIMERIC, LINKED BY ONE OR MORE DISULFIDE BONDS.
CC INHIBIN A IS A DIMER OF ALPHA AND BETA-A.
CC INHIBIN B IS A DIMER OF ALPHA AND BETA-B.
CC -!- SIMILARITY: BELONGS TO THE TGF-BETA FAMILY.
CC
CC This SWISS-PROT entry is copyright. It is produced through a collaboration
CC between the Swiss Institute of Bioinformatics and the EMBL outstation -

```

```

CC the European Bioinformatics Institute. There are no restrictions on its
CC use by non-profit institutions as long as its content is in no way
CC modified and this statement is not removed. Usage by and for commercial
CC entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC or send an email to license@isb-sib.ch).
CC -----
DR EMBL: D50327; BAA08653.1; -.
DR EMBL: U21219; AAB00874.1; -.
DR InterPro: IPR002405; Inhibin_alpha.
DR InterPro: IPR001839; TGFb.
DR Pfam: PF00019; TGF-beta; 1.
DR PRINTS: PR00669; INHIBIN.
DR PRODOM: PD000357; TGFb; 1.
DR SMART: SM00204; TGFb; 1.
DR PROSITE: PS00250; TGF_BETA_1; 1.
KW Growth factor; Hormone; Glycoprotein; Signal.
FT SIGNAL 1 20 POTENTIAL.
FT PROPEP 21 233 BY SIMILARITY.
FT CHAIN 234 367 INHIBIN ALPHA CHAIN.
FT DISULFID 263 329 BY SIMILARITY.
FT DISULFID 292 364 BY SIMILARITY.
FT DISULFID 296 366 BY SIMILARITY.
FT DISULFID 328 328 INTERCHAIN (BY SIMILARITY).
FT CARBOHYD 147 147 N-LINKED (GLCNAC. . .) (POTENTIAL).
FT CARBOHYD 269 269 N-LINKED (GLCNAC. . .) (POTENTIAL).
FT CONFLICT 105 105 Q -> R (IN REF. 2).
FT CONFLICT 171 171 G -> R (IN REF. 2).
FT CONFLICT 176 176 C -> R (IN REF. 2).
SQ SEQUENCE 367 AA; 39422 MW; DC78A6E32C84B2C61 CRC64;

Query Match 91.68; Score 131; DB 1; Length 367;
Best Local Similarity 92.08; Pred. No. 4.5e-11;
Matches 23; Conservative 1; Mismatches 1; Indels 0; Gaps 0;

QY 1 PWSPAALRLIQRPPEPSAHAFCHR 25
| | | | | | | | | | | | | | | | | | | |
DB 241 PWSPAALRLIQRPPEPSAHADCHR 265

RESULT 6
IHA_HUMAN
ID IHA_HUMAN STANDARD; PRI; 366 AA.
AC P05111;
DT 13-AUG-1987 (Rel. 05, Created)
DT 13-AUG-1987 (Rel. 05, Last sequence update)
DT 15-JUN-2002 (Rel. 41, Last annotation update)
DE Inhibin alpha chain precursor.
GN INHA.
OS Homo sapiens (Human).
OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC Mammalia; Eutheria; Primates; Catarrhini; Hominiidae; Homo.
OX NCBI_TaxId:9606;
RN [1]
RP SEQUENCE FROM N.A.
RX MEDLINE:86287450; PubMed:3016724;
RA Mayo K.E., Cerelli G.M., Spiess J., Rivier J., Rosenfeld M.G.,
RA Evans R.W., Vale W.;
RT "Inhibin A-subunit cDNAs from porcine ovary and human placenta.";
RL Proc. Natl. Acad. Sci. U.S.A. 83:5849-5853(1986).
RN [2]
RP SEQUENCE FROM N.A.
RX MEDLINE:87005283; PubMed:3758355;
RA Stewart A.G., Milborrow H.M., Ring J.M., Crowther C.E., Forage K.G.;
RT "Human inhibin genes. Genomic characterisation and sequencing.";
RL FEBS Lett. 206:329-334(1986).
RN [3]
RP SEQUENCE FROM N.A.
RC TISSUE-Brain;
RA Strausberg R.;
RL Submitted (APR-2001) to the EMBL/GenBank/DBJ databases.
RN [4]
RP SEQUENCE OF 16-366 FROM N.A.
RX MEDLINE:86186863; PubMed:3754442;

```

CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See http://www.isb-sib.ch/announce/
 CC or send an email to license@isb-sib.ch).

RA Mason A.J., Niall H.D., Seeburg P.H.;
 RT "Structure of two human ovarian inhibins";
 RL Biochem. Biophys. Res. Commun. 135:957-964(1986).
 RN [15]
 RP TISSUE SPECIFICITY, AND INDUCTION.
 RX MEDLINE: 98165602; PubMed: 9506758;
 RA Mellor S.L., Richards M.G., Pedersen J.S., Robertson D.M.,
 RA Risbridger G.P.;
 RT "Loss of the expression and localization of inhibin alpha-subunit in
 RT high grade prostate cancer";
 RL J. Clin. Endocrinol. Metab. 83:969-975(1998).
 CC -!- FUNCTION: INHIBINS AND ACTIVINS INHIBIT AND ACTIVATE.
 CC RESPECTIVELY. THE SECRETION OF FOLLITROPIN BY THE PITUITARY GLAND.
 CC INHIBINS/ACTIVINS ARE INVOLVED IN REGULATING A NUMBER OF DIVERSE
 CC FUNCTIONS SUCH AS HYPOTHALAMIC AND PITUITARY HORMONE SECRETION,
 CC GONADAL HORMONE SECRETION, GERM CELL DEVELOPMENT AND MATURATION,
 CC ERYTHROID DIFFERENTIATION, INSULIN SECRETION, NERVE CELL SURVIVAL,
 CC EMBRYONIC AXIAL DEVELOPMENT OR BONE GROWTH. DEPENDING ON THEIR
 CC SUBUNIT COMPOSITION. INHIBINS APPEAR TO OPPOSE THE FUNCTIONS OF
 CC ACTIVINS.
 CC -!- SUBUNIT: DIMERIC, LINKED BY ONE OR MORE DISULFIDE BONDS.
 CC INHIBIN A IS A DIMER OF ALPHA AND BETA-A.
 CC INHIBIN B IS A DIMER OF ALPHA AND BETA-B.
 CC -!- TISSUE SPECIFICITY: ORIGINALLY FOUND IN OVARY (GRANULOSA CELLS)
 CC AND TESTIS (SERTOLI CELLS) BUT IT IS WIDELY DISTRIBUTED IN MANY
 CC TISSUES INCLUDING BRAIN AND PLACENTA. IN ADRENAL CORTEX EXPRESSION
 CC IS LIMITED TO THE ZONA RETICULARIS AND THE INNERMOST ZONA
 CC FASCICULATA IN THE NORMAL GLAND, EXTENDING CENTRIPETALLY INTO THE
 CC ZONA FASCICULATA IN HYPERPLASIA. ALSO FOUND IN ADRENOCORTICAL
 CC TUMORS. ALSO EXPRESSED IN PROSTATE EPITHELIUM OF BENIGN PROSTATIC
 CC HYPERPLASIA. IN REGIONS OF BASAL CELL HYPERPLASIA AND IN
 CC NONMALIGNANT REGIONS OF HIGH GRADE PROSTATE CANCER. ONLY
 CC CIRCULATING INHIBIN B IS FOUND IN MALE, WHEREAS CIRCULATING
 CC INHIBINS A AND B ARE FOUND IN FEMALE.
 CC -!- SIMILARITY: BELONGS TO THE TGF-BETA FAMILY.
 CC
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC
 CC EMBL: M198; AAA59186.1; -
 CC EMBL: M1344; AAA59187.1; -
 CC EMBL: BC006391; AAH06391.1; -
 CC EMBL: X0445; CAA28040.1; -
 CC EMBL: X0446; CAA28040.1; JOINED.
 CC EMBL: A14420; CAA01158.1; -
 CC PIR: A23556; A23556.
 CC PIR: A24248; A24248.
 CC PIR: B25947; B25947.
 CC Genew: HGNC:6065; INHA.
 CC MIM: 147380; -
 CC InterPro: IPR002405; Inhibin_alpha.
 CC InterPro: IPR001839; TGFb.
 CC Pfam: PF000019; TGF-beta; 1.
 CC PRINTS: PR00669; INHIBINA.
 CC PRODOM: PD000357; TGFb; 1.
 CC SMART: SM00204; TGFb; 1.
 CC PROSITE: PS00250; TGF_BETA_1; 1.
 CC Growth factor; Hormone; Glycoprotein; Signal.
 CC SIGNAL 1 18
 CC PROPEP 19 232
 CC CHAIN 233 366
 CC DISULFID 262 328
 CC DISULFID 291 363
 CC DISULFID 295 365
 CC DISULFID 327 327
 CC CARBOHYD 146 146
 CC CARBOHYD 268 268
 CC CARBOHYD 302 302
 CC
 CC INHIBIN ALPHA CHAIN.
 CC BY SIMILARITY.
 CC BY SIMILARITY.
 CC BY SIMILARITY.
 CC INTERCHAIN (BY SIMILARITY).
 CC N-LINKED (GLCNAC. . .) (POTENTIAL).
 CC N-LINKED (GLCNAC. . .) (POTENTIAL).
 CC N-LINKED (GLCNAC. . .) (POTENTIAL).

FT CONFLICT 17 17 H -> V (IN REF. 4).
 FT CONFLICT 19 19 C -> S (IN REF. 4).
 SQ SEQUENCE 366 AA; 39669 MW; 0E03D2AB12BF8E57 CRC64;
 Query Match 89.5%; Score 128; DB 1; Length 366;
 Best Local Similarity 88.0%; Pred. No. 1.2e-10;
 Matches 22; Conservative 2; Mismatches 1; Indels 0; Gaps 0;
 QY 1 PWSPALRLRLQRPPEPSAHAFCHR 25
 DB 240 PWSPALRLRLQRPPEPSAHAFCHR 264
 |||||
 |||||
 RESULT 7
 IHA_PIG STANUARD; PRT; 364 AA.
 ID IHA_PIG
 AC P04087;
 DT 01-NOV-1986 (Rel. 03, Created)
 DT 01-APR-1988 (Rel. 07, Last sequence update)
 DT 16-OCT-2001 (Rel. 40, Last annotation update)
 DE Inhibin alpha chain precursor.
 GN INHA.
 OS Sus scrofa (Pig).
 CC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 CC Mammalia; Eutheria; Cetartiodactyla; Suina; Suidae; Sus.
 CC NCBI_TaxID=9823;
 RN [1]
 RP SEQUENCE FROM N.A.
 RX MEDLINE=86287350; PubMed=3016724;
 RA Mayo K.E., Cerelli G.M., Spiess J., Rivier J., Rosentfeld M.G.,
 RA Evans R.M., Vale W.;
 RT "Inhibin A-subunit cDNAs from porcine ovary and human placenta.";
 RL Proc. Natl. Acad. Sci. U.S.A. 83:5849-5853(1986).
 RN [2]
 RP SEQUENCE FROM N.A. AND SEQUENCE OF 231-256.
 RC TISSUE-Ovarian follicular fluid;
 RX MEDLINE=86092207; PubMed=2417121;
 RA Mason A.J., Hayflick J.S., Ling N., Esch F., Ueno N., Ying S.-Y.,
 RA Guillemin R., Niall H., Seeburg P.H.;
 RT "Complementary DNA sequences of ovarian follicular fluid inhibin show
 RT precursor structure and homology with transforming growth
 RT factor-beta";
 RL Nature 318:659-663(1985).
 CC -!- FUNCTION: INHIBINS AND ACTIVINS INHIBIT AND ACTIVATE,
 CC RESPECTIVELY. THE SECRETION OF FOLLITROPIN BY THE PITUITARY GLAND.
 CC INHIBINS/ACTIVINS ARE INVOLVED IN REGULATING A NUMBER OF DIVERSE
 CC FUNCTIONS SUCH AS HYPOTHALAMIC AND PITUITARY HORMONE SECRETION,
 CC GONADAL HORMONE SECRETION, GERM CELL DEVELOPMENT AND MATURATION,
 CC ERYTHROID DIFFERENTIATION, INSULIN SECRETION, NERVE CELL SURVIVAL,
 CC EMBRYONIC AXIAL DEVELOPMENT OR BONE GROWTH. DEPENDING ON THEIR
 CC SUBUNIT COMPOSITION. INHIBINS APPEAR TO OPPOSE THE FUNCTIONS OF
 CC ACTIVINS.
 CC -!- SUBUNIT: DIMERIC, LINKED BY ONE OR MORE DISULFIDE BONDS.
 CC INHIBIN A IS A DIMER OF ALPHA AND BETA-A.
 CC INHIBIN B IS A DIMER OF ALPHA AND BETA-B.
 CC -!- SIMILARITY: BELONGS TO THE TGF-BETA FAMILY.
 CC
 CC This SWISS-PROT entry is copyright. It is produced through a collaboration
 CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
 CC the European Bioinformatics Institute. There are no restrictions on its
 CC use by non-profit institutions as long as its content is in no way
 CC modified and this statement is not removed. Usage by and for commercial
 CC entities requires a license agreement (see <http://www.isb-sib.ch/announce/>
 CC or send an email to license@isb-sib.ch).
 CC
 CC EMBL: M1980; AAA31057.1; -
 CC EMBL: X03265; CAA27019.1; -
 CC PIR: A01392; WPPGA.
 CC PIR: A25947; A25947.
 CC InterPro: IPR002405; Inhibin_alpha.
 CC InterPro: IPR001839; TGFb.
 CC Pfam: PF000019; TGF-beta; 1.
 CC PRINTS: PR00669; INHIBINA.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

CC entries requires a license agreement (See <http://www.isb-sib.ch/announce/> or send an email to license@isb-sib.ch).

CC DR EMBL: U48438; AAA92569.1; -.
CC DR HSSP: P18075; 1BMP.
CC DR IntesPro: IPR002405; Inhibin_alpha.
CC DR InterPro: IPR001839; TGFb.
CC DR Pfam: PF00019; TGF-beta; 1.
CC DR PRINTS: P000669; INHIBINA.
CC DR PRODOM: P000357; TGFb; 1.
CC DR SMART: SM00204; TGFb; 1.
CC DR POSITE: PS00250; TGF.BETA.1; 1.
KW Growth factor; Hormone; Glycoprotein; Signal.
FT SIGNAL 1 16 POTENTIAL.
FT PROPEP 17 216
FT CHAIN 217 329 INHIBIN ALPHA CHAIN.
FT DISULFID 242 291 BY SIMILARITY.
FT DISULFID 271 326 BY SIMILARITY.
FT DISULFID 275 328 BY SIMILARITY.
FT DISULFID 290 290 INTERCHAIN (BY SIMILARITY).
FT CARBOHYD 135 135 N-LINKED (GLCNAC. . .) (POTENTIAL).
FT CARBOHYD 248 248 N-LINKED (GLCNAC. . .) (POTENTIAL).
SQ SEQUENCE 329 AA; 35853 MW; AE331687109A25A1 CRC64;

Query Match 65.0%; Score 93; DB 1; Length 329;
Best Local Similarity 68.0%; Pred. No. 8e-06;
Matches 17; Conservative 2; Mismatches 6; Indels 0; Gaps 0;

Qy 1 PWSPALRLQRPPEPSAHAFCHR 25
||||| ||||| : :||| |
Db 220 PWSPALSLLRPSDVAHTNCR 244

RESULT 10
FAS_HUMAN
ID FAS_HUMAN STANDARD; PRT; 2504 AA.
AC P49327;
DC 01-FEB-1996 (Rel. 33, Created)
DT 01-FEB-1996 (Rel. 33, Last sequence update)
DI 16-OCT-2001 (Rel. 40, Last annotation update)
DE Fatty acid synthase (EC 2.3.1.85) [Includes: EC 2.3.1.38; EC 2.3.1.39;
DE EC 2.3.1.41; EC 1.1.1.100; EC 4.2.1.61; EC 1.3.1.10; EC 3.1.2.14].
GN FASN OR FAS
OS Homo sapiens (Human).
OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
OX NCBI_TaxID=9606;
RN [1]
RP SEQUENCE FROM N.A.
RC TISSUE Brain;
RX MEDLINE-96004605; PubMed-7567999;
RA Jayakumar A., Tai M.-H., Huang W.-Y., Al-Feel W., Hsu M.,
RA Abu-Elheiga L., Chirala S.S., Wakil S.J.;
RT "Human fatty acid synthase: properties and molecular cloning";
RL Proc. Natl. Acad. Sci. U.S.A. 92:8695-8699(1995).
RN [2]
RP SEQUENCE OF 753-758 AND 1285-1297.
RX MEDLINE-94294385; PubMed-8022791;
RA Kuhajda F.P., Jenner K., Wood F.D., Hennigar R.A., Jacobs L.B.,
RA Dick J.D., Pasternack G.R.;
RT "Fatty acid synthesis: a potential selective target for
RT anticneoplastic therapy";
RL Proc. Natl. Acad. Sci. U.S.A. 91:6379-6383(1994).
CC -!- FUNCTION: FATTY ACID SYNTHASE CATALYZES THE FORMATION OF
CC LONG-CHAIN FATTY ACIDS FROM ACETYL-CoA, MALONYL-CoA AND NADPH.
CC THIS MULTIFUNCTIONAL PROTEIN HAS 7 CATALYTIC ACTIVITIES AND AN
CC ACYL CARRIER PROTEIN.
CC -!- CATALYTIC ACTIVITY: Acetyl-CoA + N malonyl-CoA + 2N NADPH = a
CC long-chain fatty acid + (N+1) CoA + N Co(2) + 2N NADP(+).
CC -!- CATALYTIC ACTIVITY: Acetyl-CoA + [acyl-carrier protein] = CoA +
CC acetyl-[acyl-carrier protein].
CC -!- CATALYTIC ACTIVITY: Malonyl-CoA + [acyl-carrier protein] = CoA +
CC malonyl-[acyl-carrier protein].

CC -!- CATALYTIC ACTIVITY: Acyl-[acyl-carrier protein] + malonyl-[acyl-
CC carrier protein] = 3-oxoacyl-[acyl-carrier protein] + Co(2) +
CC [acyl-carrier protein].
CC -!- CATALYTIC ACTIVITY: (3R)-3-hydroxyacyl-[acyl-carrier protein] +
CC NADP(+) = 3-oxoacyl-[acyl-carrier protein] + NADPH.
CC -!- CATALYTIC ACTIVITY: (3R)-3-hydroxypalmitoyl-[acyl-carrier protein]
CC = 2-hexadecenoyl-[acyl-carrier protein] + H(2)O.
CC -!- CATALYTIC ACTIVITY: Acyl-[acyl-carrier protein] + NADP(+) = trans-
CC 2,3-dehydroacyl-[acyl-carrier protein] + NADPH.
CC -!- CATALYTIC ACTIVITY: Oleoyl-[acyl-carrier protein] + H(2)O = [acyl-
CC carrier protein] + oleate.
CC -!- SUBUNIT: HOMODIMER, WHICH IS ARRANGED IN A HEAD TO TAIL FASHION.
CC -!- TISSUE SPECIFICITY: PROMINENT EXPRESSION IN BRAIN, LUNG, AND
CC LIVER.
CC -!- MISCELLANEOUS: THE RELATIVELY LOW BETA-KETOACYL SYNTHASE ACTIVITY
CC MAY BE ATTRIBUTABLE TO THE LOW PHOSPHOPANTHETHEINE CONTENT OF
CC THE PROTEIN.
CC
CC This SWISS-PROT entry is copyright. It is produced through a collaboration
CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC the European Bioinformatics Institute. There are no restrictions on its
CC use by non-profit institutions as long as its content is in no way
CC modified and this statement is not removed. Usage by and for commercial
CC entities requires a license agreement (See <http://www.isb-sib.ch/announce/>
CC or send an email to license@isb-sib.ch).

CC EMBL: U26644; AAC50259.1; -.
CC Genew; HGNC:3594; FASN.
CC MIN; 600312; .
CC InterPro: IPR001227; Acyltransferase.
CC InterPro: IPR002085; Adh_zn_family.
CC InterPro: IPR000794; Ketoacyl-synt.
CC InterPro: IPR003880; Pantne_attach.
CC InterPro: IPR000051; SAM_bind.
CC InterPro: IPR001031; Thioesterase.
CC Pfam: PF00107; adh_zinc; 1.
CC Pfam: PF00109; ketoacyl-synt; 1.
CC Pfam: PF00550; pp-binding; 1.
CC Pfam: PF00698; Acyl_transf; 1.
CC Pfam: PF00975; Thioesterase; 1.
CC Pfam: PF02601; ketoacyl-synt-C; 1.
CC PROSITE: PS00606; B-KETOACYL-SYNTHASE; 1.
CC PROSITE: PS50075; ACP_DOMAIN; 1.
KW Fatty acid biosynthesis; Multifunctional enzyme; Phosphopantetheine;
KW Hydrolase; Oxidoreductase; Transferase; Lyase; NADP;
KW Pyridoxal phosphate.
FT DOMAIN 1 413 BETA-KETOACYL SYNTHASE.
FT DOMAIN 428 815 ACYL AND MALONYL TRANSFERASES.
FT DOMAIN 1630 1857 ENOYL REDUCTASE.
FT DOMAIN 1858 2113 BETA-KETOACYL REDUCTASE.
FT DOMAIN 2118 2174 ACYL CARRIER (ACP).
FT DOMAIN 2202 2504 THIOESTERASE.
FT ACT_SITE 161 161 BETA-KETOACYL SYNTHASE (BY SIMILARITY).
FT ACT_SITE 580 580 MALONYLTRANSFERASE (BY SIMILARITY).
FT ACT_SITE 876 876 BETA-HYDROXYACYL DEHYDRATASE (BY
FT SIMILARITY).
FT NP_BIND 1666 1666 NADP (ER).
FT BINDING 1699 1699 PYRIDOXAL PHOSPHATE (BY SIMILARITY).
FT NP_BIND 1880 1895 NADP (KR).
FT BINDING 2151 2151 PHOSPHOPANTHETHEINE (BY SIMILARITY).
FT ACT_SITE 2302 2302 THIOESTERASE (BY SIMILARITY).
FT ACT_SITE 2475 2475 THIOESTERASE (BY SIMILARITY).
SQ SEQUENCE 2504 AA; 273100 MW; 8AAF9809B2738DFA CRC64;

Query Match 37.8%; Score 54; DB 1; Length 2504;
Best Local Similarity 71.4%; Pred. No. 20;
Matches 10; Conservative 2; Mismatches 2; Indels 0; Gaps 0;

Qy 9 LLORPPEPSAHAF 22
||| :|||:||||
Db 1514 LLEDKPEPTAHAF 1527

RESULTS

[illegible]

```

Query Match.      36.7%: Score 52.5; DB 1; Length 368;
Best Local Similarity 61.1%: Prod. No. 4;
Matches 11; Conservative 2; Mismatches 4; Indels 1; Gaps 1;

QY      2 WSP-AALRLIQRPPPEPS 18
      | | | | | | | | | |
Db      321 WUPRALLOCLQREPEGPA 338

```

ID	TBGL_HUMAN	STANDARD:	PRT;	451 AA.
AC	P23258; Q9BW59;			
DT	01-NOV-1991 (Rel. 20, Created)			
DT	15-JUN-2002 (Rel. 41, Last sequence update)			
DT	15-JUN-2002 (Rel. 41, Last annotation update)			
DE	Tubulin gamma-1 chain (Gamma-1 tubulin) (Gamma-tubulin complex component 1) (GCP-1).			
DE	TUBG1 OR TUBG.			
OS	Homo sapiens (Human).			
OC	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;			
OC	Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.			
OX	NCBI_TaxID=9606;			
OX	[1]			
RP	SEQUENCE FROM N.A.			
RP	MEDLINE=91249388; PubMed=1904010;			
RP	Zheng Y., Jung M.K., Oakley B.R.;			
RA	"Gamma-tubulin is present in Drosophila melanogaster and Homo sapiens and is associated with the centrosome.";			
RT	Cell 65:817-823(1991).			
RL	[2]			
RP	SEQUENCE FROM N.A.			
RP	TISSUE=Skin;			
RA	Straussberg R.;			
RL	Submitted (NOV-2000) to the EMBL/GenBank/DBJ databases.			
[3]				
RP	MASS SPECTROMETRY.			
RP	TISSUE=Breast; Cancer;			
RP	MEDLINE=21829512; PubMed=11840567;			
RP	Harris R.A., Yang A., Stein R.C., Lucy K., Brusten L., Herath A.,			
RA	Parekh R., Waterfield M.D., O'Hare M.J., Neville M.A., Page M.J.,			
RA	zvelebil M.J.;			
RT	"Cluster analysis of an extensive human breast cancer cell line			
RT	protein expression map database.";			
RT	Proteomics 2:212-223(2002).			
[4]				
RP	SUBCELLULAR LOCATION.			
RP	MEDLINE=98234404; PubMed=9566969;			
RP	Tassin A.-M., Celati C., Moudjou M., Bornet M.;			
RT	"Characterization of the human homologue of the yeast spc98p and its			
RT	association with gamma-tubulin.";			
RT	J. Cell Biol. 141:689-701(1998).			
CC	!- FUNCTION: TUBULIN IS THE MAJOR CONSTITUENT OF MICROTUBULES.			
CC	GAMMA TUBULIN IS FOUND AT MICROTUBULE ORGANIZING CENTERS (MTOC)			
CC	SUCH AS THE SPINDLE POLES OR THE CENTROSOME, SUGGESTING THAT IT			
CC	IS INVOLVED IN THE MINUS-END NUCLEATION OF MICROTUBULE ASSEMBLY.			
CC	!- SUBUNIT: Interacts with GTP2 and GCP3.			
CC	!- SUBCELLULAR LOCATION: Centrosome.			
CC	!- MASS SPECTROMETRY: MW=51197.99; METHOD=MALDI.			
CC	!- SIMILARITY: BELONGS TO THE TUBULIN FAMILY.			
CC	-----			
CC	This SWISS-PROT entry is copyright. It is produced through a collaboration			
CC	between the Swiss Institute of Bioinformatics and the EMBL outstation -			
CC	the European Bioinformatics Institute. There are no restrictions on its			
CC	use by non-profit institutions as long as its content is in no way			
CC	modified and this statement is not removed. Usage by and for commercial			
CC	entities requires a license agreement (See http://www.isb-sib.ch/			
CC	or send an email to : cense@isb-sib.ch).			
CC	-----			
DR	EMBL; M61764; AAA52620.1; -;			
DR	EMBL; BC000619; RAH00619.1; -;			
DR	PIR; A39527; UBUU0.			
DR	GeneW; HGNC:12417; TUBG1.			
DR	MIM; 191135; -;			
DR	InterPro; IPR000217; Tubulin.			
DR	InterPro; IPR003008; Tubulin_FtsZ.			
DR	Pfam; PF00091; tubulin; 1.			
DR	PRINTS; PR01161; TUBULIN.			
DR	PROSITE; PS00227; TUBULIN; 1.			
DR	Microtubules; GTP-binding.			
FT	NP_BIND 142 148 GTP (POTENTIAL).			

FT CONFLICT 35 35 G -> A (IN REF. 1).
ST CONFLICT 202 202 V -> L (IN REF. 1).
SQ SEQUENCE 451 AA: 51170 MW: E2A4C0179ED0CFE8 CRC64;

Query Match: 35.7%; Score 51; DB 1; Length 451;
Best Local Similarity 45.5%; Pred. No. 8.2;
Matches 10; Conservative 5; Mismatches 5; Indels 2; Gaps 1;

QY 1 PWSPAALR--LLQRPPEPSAH 20

|||||

Db 350 PWGPASIQVALSRKSPYLPSAH 371

RESULT 13

TBG1_MOUSE
ID TBG1_MOUSE STANDARD; PRI; 451 AA.
AC O9Z310;
DT 15-JUN-2002 (Rel. 41, Created)
DT 15-JUN-2002 (Rel. 41, Last sequence update)
DT 15-JUN-2002 (Rel. 41, Last annotation update)
DE Tubulin gamma-1 chain (Gamma-1 tubulin) (Gamma-tubulin complex component 1) (GCP-1).
GN TBG1 OR TUBG.
OS Mus musculus (Mouse), and
OS Rattus norvegicus (Rat).
OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Mus.
OX NCBI_TaxID=10090, 10116;
RN [1]
RP SEQUENCE FROM N.A.
RC SPECIES-Mouse;
RA Strausberg K.;
RL Submitted (APR-2001) to the EMBL/GenBank/DDBJ databases.
RT [2]
RN [2]
RP SEQUENCE FROM N.A.
RC SPECIES Rat;
RX MEDLINE=99397453; PubMed=10470852;
RA Nakadai T., Okada N., Makino Y., Tamura T.;
RI "Structure of rat gamma-tubulin and its binding to HP33.";
RL DNA Res. 6:207-209(1999).
CC -!- FUNCTION: TUBULIN IS THE MAJOR CONSTITUENT OF MICROTUBULES. IT BINDS TWO MOLES OF GTP, ONE AT AN EXCHANGEABLE SITE ON THE BETA CHAIN AND ONE AT A NONEXCHANGEABLE SITE ON THE ALPHA CHAIN (BY SIMILARITY).
CC -!- SUBUNIT: Interacts with GCP2 and GCP3 (By similarity).
CC -!- SUBCELLULAR LOCATION: Centrosome (By similarity).
CC -!- SIMILARITY: BELONGS TO THE TUBULIN FAMILY.

CC This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See <http://www.isb-sib.ch/announce/> or send an email to license@isb-sib.ch).

CC EMBL; W006581; AAR0581.1;
CC EMBL; AB015946; BAA36504.1;
CC MGD; MGI:101834; Tuba9.

DR InterPro; IPR000217; Tubulin.
DR InterPro; IPR003008; Tubulin_FtsZ.
DR Pfam; PF00091; tubulin; 1.
DR PRINTS; PR01161; TUBULIN.

DR PROSITE; PS00227; TUBULIN; 1.

KW Microtubules; GTP-binding.

FT NP_BIND 142 148 GTP (POTENTIAL).

SQ SEQUENCE 451 AA: 51100 MW: A8F106BD12D0C88A CRC64;

Query Match: 35.7%; Score 51; DB 1; Length 451;
Best Local Similarity 45.5%; Pred. No. 8.2;
Matches 10; Conservative 5; Mismatches 5; Indels 2; Gaps 1;

QY 1 PWSPAALR--LLQRPPEPSAH 20

|||||

Db 350 PWGPASIQVALSRKSPYLPSAH 371

RESULT 14

TBG2_HUMAN
ID TBG2_HUMAN STANDARD; PRI; 451 AA.
AC Q9NRH3;
DT 16-OCT-2001 (Rel. 40, Created)
DT 16-OCT-2001 (Rel. 40, Last sequence update)
DT 15-JUN-2002 (Rel. 41, Last annotation update)
DE tubulin gamma-2 chain (Gamma-2 tubulin).
GN TUBG2.
OS Homo sapiens (Human).
OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
OC Mammalia; Eutheria; Primates; Catartini; Hominidae; Homo.
OX NCBI_TaxID=9606;
RN [1]
RP SEQUENCE FROM N.A.
RX MEDLINE=20366133; PubMed=10903841;
RI Wise D.O., Krahe R., Oakley B.R.;
RT "The gamma-tubulin gene family in humans.";
RL Genomics 87:164-170(2000).
RN [2]
RP SEQUENCE FROM N.A.
RC TISSUE=Mammary gland;
RA Isogai T., Ota T., Hayashi K., Sujiyama T., Otsuki T., Suzuki Y., Nishikawa T., Nagai K., Sugano S., Shiratori A., Sudo H., Nagatsuna M., Hosoi T., Kaku Y., Kodaira H., Kondo H., Sugawara M., Takahashi M., Chiba Y., Ishida S., Murakawa K., Ono Y., Takiguchi S., Matarabe S., Kimura K., Murakami K., Ishii S., Kawai Y., Saito K., Yamamoto J., Wakamatsu A., Nakamura Y., Nagahari K., Masuho Y., Ninomiya K., Iwayanagi T.;
RT "NEO human cDNA sequencing project.";
RL Submitted (AUG-2000) to the EMBL/GenBank/DDBJ databases.
RN [3]
RP SEQUENCE FROM N.A.
RC TISSUE=Lung;
RA Strausberg K.;
RL Submitted (JUN-2001) to the EMBL/GenBank/DDBJ databases.
CC -!- FUNCTION: TUBULIN IS THE MAJOR CONSTITUENT OF MICROTUBULES. GAMMA TUBULIN IS FOUND AT MICROTUBULE ORGANIZING CENTERS (MTOC) SUCH AS THE SPINDLE POLES OR THE CENTROSOME, SUGGESTING THAT IT IS INVOLVED IN THE MINUS-END NUCLEATION OF MICROTUBULE ASSEMBLY.
CC -!- SIMILARITY: BELONGS TO THE TUBULIN FAMILY.

CC This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See <http://www.isb-sib.ch/announce/> or send an email to license@isb-sib.ch).

CC EMBL; AF225971; AAF34188.1;
CC EMBL; AK022324; BAB14012.1;
CC EMBL; BC009670; AAH09670.1;
CC Genew; HGNC:12419; TUBG2.

CC MIM; 605785;

DR InterPro; IPR000217; Tubulin.

DR InterPro; IPR003008; Tubulin_FtsZ.

DR Pfam; PF00091; tubulin; 1.

DR PRINTS; PR01161; TUBULIN.

DR PROSITE; PS00227; TUBULIN; 1.

KW Microtubules; GTP-binding.

FT NP_BIND 142 148 GTP (POTENTIAL).

SQ SEQUENCE 451 AA: 51091 MW: FDE2CB5A33D92691 CRC64;

Query Match: 35.7%; Score 51; DB 1; Length 451;
Best Local Similarity 45.5%; Pred. No. 8.2;
Matches 10; Conservative 5; Mismatches 5; Indels 2; Gaps 1;

QY 1 PWSPAALR--LLQRPPEPSAH 20

```

Db 350 FWGPASIQVALSRKSPYLP5AH 371
||||:|:| ||||
RESULT 15
TBG2_MOUSE
ID TBG2_MOUSE STANDARD; PRT: 451 AA.
AC Q8VCK3;
DT 15-JUN-2002 (rel. 41, Created)
DT 15-JUN-2002 (rel. 41, Last sequence update)
DT 15-JUN-2002 (rel. 41, Last annotation update)
DE Tubulin gamma-2 chain (Gamma-2 tubulin).
GN TBG2.
OS Mus musculus (Mouse).
OC Eukaryota; Metazoa; Chordata; Vertebrata; Euteleostomi;
OC Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Mus.
OX NCBI_TaxID=10090;
RN [1]
RP SEQUENCE FROM N.A.
RC TISSUE=Retina;
RA Strausberg R.;
RL Submitted (DEC-2001) to the EMBL/GenBank/DDBJ databases.
CC -!- FUNCTION: TUBULIN IS THE MAJOR CONSTITUENT OF MICROTUBULES.
CC GAMMA TUBULIN IS FOUND AT MICROTUBULE ORGANIZING CENTERS (MTOC)
CC SUCH AS THE SPINDLE POLES OR THE CENTROSOME, SUGGESTING THAT IT
CC IS INVOLVED IN THE MINUS-END NUCLEATION OF MICROTUBULE ASSEMBLY.
CC -!- SIMILARITY: BELONGS TO THE TUBULIN FAMILY.
CC -----
CC THIS SWISS-PROT entry is copyright. It is produced through a collaboration
CC between the Swiss Institute of Bioinformatics and the EMBL outstation -
CC the European Bioinformatics Institute. There are no restrictions on its
CC use by non-profit institutions as long as its content is in no way
CC modified and this statement is not removed. Usage by and for commercial
CC entities requires a license agreement (See http://www.isb-sib.ch/announce/
CC or send an email to license@isb-sib.ch).
CC -----
DR EMBL: BF019652; AAB19652.1; -.
DR InterPro: IPR000217; Tubulin.
DR InterPro: IPR003008; Tubulin_FtsZ.
DR PROSITE: PS00227; TUBULIN; 1.
KW Microtubules; GTP-binding.
FT NP_BIND 142 148 GTP (POTENTIAL).
SQ SEQUENCE 451 AA; 51121 MW; 1FCC0FE1E04839AA CRC64;

Query Match 35.7%; Score 51; DR 1; Length 451;
Best local Similarity 45.5%; Pred. No. 8.2;
Matches 10; Conservative 5; Mismatches 5; Indels 2; Gaps 1;

QY 1 FWGPASLR-LIQRPPEPSAH 20
||||:|:| ||||
Db 350 FWGPASIQVALSRKSPYLP5AH 371

Search completed: March 13, 2003, 12:35:43
Job time : 13 secs

```